

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

Pearson Edexcel Level 3 GCE**Friday 7 June 2024**

Afternoon (Time: 1 hour 30 minutes)

Paper
reference**9FM0/3C****Further Mathematics****Advanced****PAPER 3C: Further Mechanics 1****You must have:**

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P75322A

©2024 Pearson Education Ltd.
F:1/1/

Pearson

1. [In this question, $\mathbf{i}$ and $\mathbf{j}$ are horizontal perpendicular unit vectors.]

A particle A has mass 3 kg and a particle B has mass 2 kg.

The particles are moving on a smooth horizontal plane when they collide directly.

Immediately **before** the collision, the velocity of A is $(3\mathbf{i} - \mathbf{j})\text{ms}^{-1}$ and the velocity of B is $(-6\mathbf{i} + 2\mathbf{j})\text{ms}^{-1}$

Immediately **after** the collision the velocity of A is $\left(-2\mathbf{i} + \frac{2}{3}\mathbf{j}\right)\text{ms}^{-1}$

- (a) Find the total kinetic energy of the two particles **before** the collision. (3)
- (b) Find, in terms of $\mathbf{i}$ and $\mathbf{j}$, the impulse exerted on A by B in the collision. (3)
- (c) Find, in terms of $\mathbf{i}$ and $\mathbf{j}$, the velocity of B immediately **after** the collision. (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 is 9 marks)

2. A rough plane is inclined to the horizontal at an angle θ , where $\tan \theta = \frac{3}{4}$

A particle P of mass m is at rest at a point on the plane.

The particle is projected **up** the plane with speed $\sqrt{2ag}$

The particle moves up a line of greatest slope of the plane and comes to instantaneous rest after moving a distance d .

The coefficient of friction between P and the plane is $\frac{1}{7}$

- (a) Show that the magnitude of the frictional force acting on P as it moves up the plane

is $\frac{4mg}{35}$ (3)

Air resistance is assumed to be negligible.

Using the work-energy principle,

- (b) find d in terms of a . (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 2 is 7 marks)



This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 3 is 12 marks)

4. A particle A of mass $2m$ is moving in a straight line with speed $3u$ on a smooth horizontal plane. Particle A collides directly with a particle B of mass m which is at rest on the plane.

The coefficient of restitution between A and B is e , where $e > 0$

- (a) Show that the speed of B immediately after the collision is $2u(1 + e)$. (6)

After the collision, B hits a smooth fixed vertical wall which is perpendicular to the direction of motion of B .

- (b) Show that there will be a second collision between A and B . (3)

The coefficient of restitution between B and the wall is $\frac{1}{2}$

Find, **in simplified form**, in terms of m , u and e ,

- (c) the magnitude of the impulse received by B in its collision with the wall, (3)

- (d) the loss in kinetic energy of B due to its collision with the wall. (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 is 15 marks)

5. A light elastic string has natural length $2a$ and modulus of elasticity $2mg$. One end of the string is attached to a fixed point A on a horizontal ceiling. The other end is attached to a particle P of mass m .

The particle P hangs in equilibrium at the point E , where $AE = 3a$.

The particle P is then projected vertically downwards from E with speed $\frac{3}{2}\sqrt{ag}$

Air resistance is assumed to be negligible.

Find the elastic energy stored in the string, when P first comes to instantaneous rest. Give your answer in the form $kmg a$, where k is a constant to be found.

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 5 is 7 marks)

P 7 5 3 2 2 A 0 1 5 2 4

6. [In this question, $\mathbf{i}$ and $\mathbf{j}$ are horizontal perpendicular unit vectors.]

A particle P is moving with velocity $(4\mathbf{i} - \mathbf{j})\text{ m s}^{-1}$ on a smooth horizontal plane.
The particle collides with a smooth vertical wall and rebounds with velocity $(\mathbf{i} + 3\mathbf{j})\text{ m s}^{-1}$

The coefficient of restitution between P and the wall is e .

(a) Find the value of e .

(6)

After the collision, P goes on to hit a second smooth vertical wall, which is parallel to $\mathbf{i}$.

The coefficient of restitution between P and this second wall is $\frac{1}{3}$

The angle through which the direction of motion of P has been deflected by its collision with this second wall is α° .

(b) Find the value of α , giving your answer to the nearest whole number.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 6 is 10 marks)

7.




Figure 1

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

A smooth uniform sphere A of mass m is moving with speed U on a smooth horizontal plane. The sphere A collides obliquely with a smooth uniform sphere B of mass $3m$ which is at rest on the plane. The two spheres have the same radius.

Immediately before the collision, the direction of motion of A makes an angle α , where $0^\circ < \alpha < 90^\circ$, with the line joining the centres of the spheres.

Immediately after the collision, the direction of motion of A is **perpendicular** to its original direction, as shown in Figure 1.

The coefficient of restitution between the spheres is e .

(a) Show that the speed of B immediately after the collision is

$$\frac{1}{4}(1+e)U\cos\alpha \quad (6)$$

(b) Show that $e > \frac{1}{3}$ (4)

(c) Show that $0 < \tan\alpha \leq \frac{1}{\sqrt{2}}$ (5)



[illegible]

DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 7 is 15 marks)

TOTAL FOR PAPER IS 75 MARKS

